

Title The role of volatile compound in recognition and germination of *Penicillium digitatum* and *Penicillium italicum* on citrus fruit

Author Droby S, Eick A, Macarasin D, Cohen L and Rafael G

Citation Program and Abstracts, 11th International Citrus Congress (ISC Congress), 26-20 October 2008, Wuhan, China. 333 pages.

Keyword citrus; volatile compound; green mould; blue mould

Abstract

Volatiles emitted from wounded peel tissue of various citrus cultivars had a pronounced stimulatory effect on germination and germ tube elongation of both *P. digitatum* and *P. italicum*; however, *P. digitatum* appeared to be more sensitive to the stimulatory action of citrus peel volatiles. When exposed to volatiles from grapefruit, the percentage of germinated spores of *P. digitatum* and *P. italicum* was 10 and 5 fold, respectively as compared to the control. In contrast, *B. cinerea* and *P. expansum* were either not affected or inhibited by the peel volatiles. GS-MS analysis of volatiles present in the peel of various citrus fruit cultivars revealed that limonene is the major fruit peel volatile. Its percentage ranged from 89-95% at the early stages of fruit development throughout the harvest season. Myrcene and α -pinene made up the second and third greatest amounts among the volatiles found in these oils. All four monoterpenes, limonene α -pinene, β -pinene and myrcene were stimulatory to *P. digitatum* and *P. italicum* but inhibitory to or had no effect on *P. expansum* and *B. cinerea*. Germ tube elongation in *P. digitatum* responded most strongly to limonene and less strongly to α -pinene and β -pinene while myrcene had little effect. In *P. italicum* myrcene stimulated germ tube elongation the most followed by limonene, with α -pinene, and β -pinene being about equal. Germination of *P. italicum* conidia was highest in response to myrcene with the effect of the other compounds being about equal at concentrations of 5 μ l or more per plate.